

The Environmental and Health Effects of the Coal Ash Spill at Kingston, Tennessee: Preliminary assessment

Avner Vengosh¹, Laura Ruhl¹, Gary S. Dwyer¹, Heileen Hsu-Kim², Amrika Deonarine², Mike Bergin³, and Julia Kravchenko⁴

¹Division of Earth and Ocean Sciences, Nicholas School of the Environment, 205 Old Chemistry Building, Box 90227, Duke University, Durham, NC 27708; ²Civil and Environmental Engineering, 121 Hudson Hall, Box 90287, Duke University, Durham, NC 27708; ³School of Civil and Environmental Engineering and School of Earth and Atmospheric Sciences, 311 Ferst Drive, Georgia Institute of Technology, Atlanta, GA, 30332; ⁴ Duke Comprehensive Cancer Center in Duke School of Medicine, 2424 Erwin Road, Hock Plaza, Suite 601, Durham, NC 27705.

KEYWORDS: TVA coal ash spill, water contamination, trace metals, arsenic, radium, environment, health

ABSTRACT

A preliminary investigation of the potential environmental and health effects of the largest coal ash spill in USA history at Tennessee Valley Authority (TVA) Kingston coal-burning power plant has revealed three major effects. First, the surficial release of coal ash formed a sub-aerial deposit that contains high levels of toxic elements (As=75 mg/kg; Hg= 150 mg/kg) and radioactivity ($^{226}\text{Ra}+^{228}\text{Ra}=8$ pCi/g). These pose a potential health risk to local communities as a possible source of air-borne resuspended fine particles (<10 μm). Second, high concentrations of contaminants from the coal ash leachate in the aquatic environments have caused severe water contamination in areas of restricted water exchange. Farther downstream, in the Emory and Clinch Rivers, much lower levels of these elements were found due to river dilution. Third, accumulation and re-mobilization of Hg- and As-rich coal ash in river sediments have the potential to impact aquatic ecosystems in the downstream rivers, especially by methyl-mercury formation in anaerobic river sediments.